



Post Operative Care

From your preeminent Medical Concierge in Antalya



BREAST ENLARGEMENT

www.myimperialcare.com

BREAST ENLARGEMENT

WEARING YOUR BRA

A bra will be placed on you at the time of your surgery. If your bra feels too tight or hurts, it may cause ulceration. To avoid this, you may unhook the bottom two hooks to make it more comfortable. For tissue healing, it is necessary to wear a bra at all times for four weeks. You may remove the bra for showering only. If you have incisions in the crease of your breast, do not wear a bra with an underwire until sensation to the scar has returned to normal, with full sensation.

STERI-STRIPS

Steri-strips are adhesive strips that support the incisions in their early stages of healing and prevent widening of scars. Please leave the steri-strips on until they begin to come off naturally. You may gently wash over the steri-strips with anti-bacterial soap. Do not trim or pull any suture. The long tails of the sutures are needed when removing them.

BLEEDING

Small amounts of oozing and bleeding are common and expected. Feel free to change the gauze as needed at home. If the bleeding is more than a slow staining of the dressings, apply firm pressure for 20-30 minutes. Should heavy bleeding occur, apply firm pressure and call us immediately at +90 542 512 51 64.

BRUISING AND SWELLING

Bruising and swelling are to be expected. Swelling will take 3-6 months to fully resolve. The shape of your breasts will change and appear more natural and relaxed over several months.

SHOWERING AND BATHING

You may take a shower 24 hours after surgery. Strictly avoid tub baths, hot tubs, swimming pools, oceans, and lakes until the incision is completely healed and no longer forming scabs.



PAIN

Mild to moderate discomfort is normal after surgery. Please follow instructions given to you with your pain medications. You may hear and feel fluid swooshing around your implant, this is normal.

DRIVING

You may drive when driving does not cause you pain. You must wait 24 hours after anesthesia and until you are no longer taking pain medications before you resume driving. This would be driving under the influence and you would endanger yourself and others.

HEMATOMAS

Hematomas are the collection of blood and fluid under the skin or muscle. They are rare, but must be reported immediately. A hematoma is evidenced by one breast becoming significantly more swollen and painful than the other breast. If you see evidence of a hematoma, contact us immediately or visit the nearest hospital (if you have traveled back to your home country) to be assessed and possibly let them drain the hematoma.

ACTIVITY

Do not lift anything over 5 pounds for 2-3 days. As you begin to resume normal activity, refrain from high impact or strenuous activities, such as weight lifting, upper body workout, and jogging for 4 weeks. You may resume light walking as tolerated. You should discuss your exercise routine with us so that we can help you with proper guidance.

INFECTION

Although infection is unusual after this type of surgery, please take all prescribed antibiotics until full prescriptions is empty. Signs of infection include drainage with a foul odor, yellow or green drainage from the incision, fever of 101 degrees (usually accompanied by chills), or redness and warmth around the incision. These are symptoms which need to be reported. Smokers have a greater risk of skin loss and wound healing complications. Do not smoke while your body is trying to heal from surgery.

